



ELSEVIER

Journal of Computational and Applied Mathematics 51 (1994) 179–191

---



---

JOURNAL OF  
COMPUTATIONAL AND  
APPLIED MATHEMATICS

---



---

# Rates of convergence for collocation with Jacobi polynomials for the airfoil equation

T. Hartmann<sup>1</sup>, E.P. Stephan<sup>\*,2</sup>

*Institut für Angewandte Mathematik, Universität Hannover, Welfengarten 1, D-30167 Hannover, Germany*

Received 23 April 1992; revised 14 August 1992

---

## Abstract

In this paper we consider for the classical airfoil equation a collocation method based on Jacobi polynomials. The error is estimated in suitable nonstandard Sobolev norms which are defined in such a way to respect the singularity structure of the exact solution. Furthermore, our numerical experiments underline our optimal theoretical error estimates.

**Key words:** Collocation method; Airfoil equations; Jacobi polynomials

---

## 0. Introduction

The classical airfoil equation

$$Sv(x) := \frac{1}{\pi} \int_{-1}^1 \frac{v(y)}{y-x} dy = f(x), \quad -1 < x < 1, \quad (1)$$

has for smooth  $f$  a solution of the form

$$v(x) = \rho(x)u(x),$$

where  $\rho(x)$  is the weight function

$$\rho(x) = (1-x)^\alpha (1+x)^\beta \quad (2)$$

and  $u(x)$  is “smooth”. Here the integral is understood as a Cauchy principal value integral.

---

\* Corresponding author.

<sup>1</sup> Supported by DFG grant Ste 573/1-1.

<sup>2</sup> Partially supported by the University of New South Wales Australian Research Council program grant “Numerical analysis for integrals, integral equations and boundary value problems” when visiting the School of Mathematics of the University of New South Wales, Sydney, Australia. The author also gratefully acknowledges the support of the Center for Mathematical Analysis of the Australian National University, Canberra, Australia.

The collocation method for the above equation to be considered here based on Chebyshev (Jacobi) polynomials is perhaps the easiest method of obtaining a numerical solution. It correctly represents the endpoint singularities of the exact solution, and yields faster than polynomial convergence if  $f$  is smooth.

We do not claim that the method is new. It has, for example, been suggested by Golberg [9,10] and others [1,3–6,11]. Our emphasis in the present work is on proving optimal rates of convergence in suitable Sobolev spaces (Theorems 1 and 6), by means of an analysis closely related to that used in [7,8,13]. Our analysis in Sections 1 and 2 is based on various discrete inner products. Numerical experiments which underline our theoretical results are given in Section 3.

In the following we list some well-known facts on Jacobi polynomials and explicit mapping properties of the integral operator  $S$ .

Let  $\{p_j^{(\alpha,\beta)}(x)\}$ ,  $j \geq 0$ , be the Jacobi polynomials orthogonal with respect to the weight function  $\rho(x)$ , i.e.,  $\{p_n^{(1/2,-1/2)}(x)\}_0^\infty$ ,  $\{p_n^{(-1/2,1/2)}(x)\}_0^\infty$  are Jacobi polynomials, normalized by the conditions

$$\int_{-1}^1 (1-x)^{1/2}(1+x)^{-1/2} p_n^{(1/2,-1/2)}(x) p_m^{(1/2,-1/2)}(x) dx = \delta_{mn},$$

$$\int_{-1}^1 (1+x)^{1/2}(1-x)^{-1/2} p_n^{(-1/2,1/2)}(x) p_m^{(-1/2,1/2)}(x) dx = \delta_{mn}.$$

$\{p_n^{(1/2,-1/2)}(x)\}_0^\infty$  is a complete orthonormal basis for  $L_{2,\rho}$ ,  $\rho = (1-x)^{1/2}(1+x)^{-1/2}$ ,  $\{p_n^{(-1/2,1/2)}(x)\}_0^\infty$  is a complete orthonormal basis for  $L_{2,1/\rho}$ . Here  $\alpha$  and  $\beta$  determined by

$$\alpha = \frac{1}{2} + m, \quad \beta = -\frac{1}{2} + n, \quad (3)$$

and the integers  $m, n$  are related to the index  $\nu$  of (1) by

$$\nu = -(\alpha + \beta) = -(m + n).$$

We will consider only nonnegative values of the index and, in order to have integrable solutions, we take  $\alpha, \beta > -1$ . This restricts the index  $\nu$  and the exponents  $\alpha, \beta$  to the cases

$$\begin{aligned} (1) \quad & \alpha = -\frac{1}{2}, \quad \beta = -\frac{1}{2}, \quad \nu = 1, \\ (2) \quad & \alpha = \frac{1}{2}, \quad \beta = -\frac{1}{2}, \quad \nu = 0, \\ (3) \quad & \alpha = -\frac{1}{2}, \quad \beta = \frac{1}{2}, \quad \nu = 0. \end{aligned} \quad (4)$$

When  $\nu = 0$ , the solution of (1) is unique whereas when  $\nu = 1$ , equation (1) usually has an eigensolution; to find a unique solution an additional side condition must be imposed like

$$l(u) = c, \quad (5)$$

where  $l$  is a given linear functional and  $c$  is a given constant.

Next, we introduce the notation for  $\rho(x)$  from (2) with exponents as in (3):

$$S_\rho u(x) = \frac{1}{\pi} \int_{-1}^1 \frac{\rho(y)u(y)}{y-x} dy.$$

When  $\nu = 0$ , there holds (see [9])

$$S_\rho p_j^{(\alpha, \beta)}(x) := \frac{-1}{\sin(\pi\alpha)} p_j^{(-\alpha, -\beta)}(x), \quad j = 0, 1, \dots,$$

i.e., for  $\rho = (1-x)^{1/2}(1+x)^{-1/2}$ ,

$$S_\rho p_j^{(1/2, -1/2)}(x) = -p_j^{(-1/2, 1/2)}(x), \quad j = 0, 1, \dots, \quad (6)$$

and for  $\rho = (1-x)^{-1/2}(1+x)^{1/2}$ ,

$$S_\rho p_j^{(-1/2, 1/2)}(x) = p_j^{(1/2, -1/2)}(x), \quad j = 0, 1, \dots$$

When  $\nu = 1$ , there holds (see [9])

$$S_\rho p_j^{(\alpha, \beta)}(x) = \begin{cases} -\frac{1}{\sin(\pi\alpha)} p_{j-1}^{(-\alpha, -\beta)}(x), & j = 1, 2, \dots, \\ 0, & j = 0, \end{cases}$$

i.e., for  $\rho = (1-x)^{-1/2}(1+x)^{-1/2}$ ,

$$S_\rho p_j^{(-1/2, -1/2)}(x) = \begin{cases} p_{j-1}^{(1/2, 1/2)}(x), & j = 1, 2, \dots, \\ 0, & j = 0. \end{cases} \quad (7)$$

The case (3) in (4) does not inherit any substantial differences to case (2), so we consider only the cases (1) and (2).

## 1. Method A

In order to make the paper more readable we use for the Method A, i.e.,  $\alpha = \beta = -\frac{1}{2}$ ,  $\nu = 1$ , the common notation for Chebyshev polynomials. There holds from (7),

$$S_\rho T_j = U_{j-1}, \quad j = 1, 2, \dots, \quad S_\rho T_0 = 0,$$

where  $T_j = p_j^{(-1/2, -1/2)}$  are the Chebyshev polynomials of the first kind and  $U_j = p_j^{(1/2, 1/2)}$  are the Chebyshev polynomials of the second kind normalized by

$$\int_{-1}^1 (1-x^2)^{-1/2} T_j(x) T_k(x) dx = \begin{cases} \delta_{jk} \cdot \frac{1}{2}\pi, & k \neq 0, \\ \delta_{jk} \cdot \pi, & k = 0, \end{cases}$$

and

$$\int_{-1}^1 (1-x^2)^{1/2} U_j(x) U_k(x) dx = \delta_{jk} \cdot \frac{1}{2}\pi.$$

We approximate the solution  $\nu$  of (1) by  $\rho(x) = 1/\sqrt{1-x^2}$  times a polynomial  $u_n$  of degree  $\leq n-1$ :

$$v_n(x) = \rho(x)u_n(x) = \rho(x) \left( \sum_{k=0}^{n-1} a_k T_k(x) \right), \quad a_k \in \mathbb{R}, \quad k = 0, \dots, n-1. \quad (8)$$

We choose for the collocation points

$$\tilde{x}_j = \cos \frac{j}{n} \pi, \quad 1 \leq j \leq n-1,$$

the  $n-1$  zeros of  $U_{n-1}(x)$ . Thus our collocation scheme is: find  $u_n \in \mathbb{P}_{n-1}$ , such that

$$S_\rho u_n(\tilde{x}_j) = f(\tilde{x}_j), \quad 1 \leq j \leq n-1. \quad (9)$$

Here  $\mathbb{P}_{n-1}$  denotes the space of polynomials of degree  $\leq n-1$  on  $(-1, 1)$ . Now, with the aid of (7), we have

$$S_\rho u_n(x) = \sum_{k=1}^{n-1} a_k U_{k-1}(x). \quad (10)$$

On the other hand, the choice  $c = 0$  and

$$l(u_n) = \int_{-1}^1 \rho(x) u_n(x) dx \stackrel{!}{=} 0 \quad (11)$$

in (5) leads to the constraint

$$a_0 = 0.$$

Thus equations (9) and (11) represent a set of linear equations for  $a_0, a_1, \dots, a_{n-1}$  with easily computed matrix elements. One may even obtain explicit expressions for  $a_1, \dots, a_{n-1}$  by exploiting the discrete orthogonality of Chebyshev polynomials of the second kind. Let us define an inner product incorporating the weight  $1/\rho$  by

$$\langle\langle u, w \rangle\rangle = \int_{-1}^1 u(x) w(x) \frac{1}{\rho(x)} dx, \quad (12)$$

and a corresponding discrete inner product (obtained by using the Gaussian quadrature rule for the weight  $(1-x)^{1/2}(1+x)^{1/2}$  where the nodes  $x_k$  are the zeros of  $U_n(x)$ )

$$\langle\langle u, w \rangle\rangle_n = \sum_{k=1}^n \omega_k u(x_k) w(x_k).$$

We have

$$\omega_k = \frac{\pi}{n+1} \sin^2 \left( \frac{k}{n+1} \pi \right), \quad k = 1, \dots, n.$$

Then, as the  $n$ -point Gaussian quadrature rule is exact for all polynomials of degree  $\leq 2n-1$ , we have, for  $j, k = 0, \dots, n-1$ , the discrete orthogonality property

$$\langle\langle U_j, U_k \rangle\rangle_n = \langle\langle U_j, U_k \rangle\rangle = \delta_{jk} \cdot \frac{1}{2} \pi. \quad (13)$$

It then follows from (9), (10) and (13) that the coefficients in (8) have the explicit form

$$a_j = \frac{2}{\pi} \langle\langle f, U_{j-1} \rangle\rangle_{n-1}, \quad j = 1, \dots, n-1, \quad a_0 = 0.$$

In particular, if  $f \in \mathbb{P}_{n-1}$ , then Method A yields the exact solution.

In order to present the convergence results, we begin by defining some norms by which the error may be described.

Let  $u$  have a Chebyshev polynomial expansion

$$u(x) = \sum_{k=0}^{\infty} \tilde{u}(k) T_k(x), \quad \text{with} \quad \begin{cases} \tilde{u}(k) = \frac{2}{\pi} \langle u, T_k \rangle, & k \in \mathbb{Z}_+, \\ \tilde{u}(0) = \frac{1}{\pi} \langle u, T_0 \rangle, \end{cases} \quad (14)$$

and  $\tilde{u}(0) = 0$  by (11) where

$$\langle u, w \rangle = \int_{-1}^1 \rho(x) u(x) w(x) dx.$$

Then we define the space  $\tilde{H}_\rho^s$ ,  $s \in \mathbb{R}$ , to be the closure of all polynomials with respect to the norm

$$\|u\|_{\tilde{H}_\rho^s}^2 = \sum_{k=1}^{\infty} |k|^{2s} |\tilde{u}(k)|^2 + |\tilde{u}(0)|^2.$$

Note  $\|u\|_{\tilde{H}_\rho^0}^2 = \langle u, u \rangle = \|u\|_{L_{2,\rho}}^2$ .

**Theorem 1.** Let  $u \in \tilde{H}_\rho^t$  solve (1) with  $u = v/\rho$  and  $\rho(x) = (1-x^2)^{-1/2}$ . Then for any  $n \in \mathbb{Z}_+$  there exists a solution  $u_n \in \mathbb{P}_{n-1}$  of (9). Furthermore, if  $t > s$  and  $t > \frac{1}{2}$ , then  $u_n$  satisfies

$$\|u_n - u\|_{\tilde{H}_\rho^s} \leq \begin{cases} C n^{-(t-s)} \|u\|_{\tilde{H}_\rho^t}, & s \geq 0, \\ C n^{-t} \|u\|_{\tilde{H}_\rho^t}, & s < 0, \end{cases}$$

with  $C$  a constant independent of  $n$ .

**Remark 2.** In [9] Golberg shows the convergence of the above methods with a different analysis, but there error estimates are not given.

**Theorem 3.** The operator  $S_\rho: \tilde{H}_\rho^s \setminus \{T_0\} \rightarrow \hat{H}_{1/\rho}^s$  is continuous with continuous inverse where the space  $\hat{H}_{1/\rho}^s$ ,  $s \in \mathbb{R}$ , is the closure of all polynomials with respect to the norm

$$\|f\|_{\hat{H}_{1/\rho}^s} = \left( \sum_{k=0}^{\infty} |k+1|^{2s} |\hat{f}(k)|^2 \right)^{1/2},$$

where  $\hat{f}(k) = 2/\pi \langle f, U_k \rangle$ ,  $k \in \mathbb{Z}_+$ , with (12).

**Proof of Theorem 1.** The approximation scheme (9) is equivalent to

$$\langle \langle S_\rho u_n, U_j \rangle \rangle_{n-1} = \langle \langle S_\rho u, U_j \rangle \rangle_{n-1}, \quad j = 0, \dots, n-2. \quad (15)$$

Now, from (7) and (14) we have

$$S_\rho u(x) = \sum_{k=1}^{\infty} \tilde{u}(k) U_{k-1}(x).$$

Thus,

$$\langle\langle S_\rho u, U_j \rangle\rangle_{n-1} = \sum_{k=1}^{\infty} \tilde{u}(k) \langle\langle U_{k-1}, U_j \rangle\rangle_{n-1}, \quad j = 0, \dots, n-2,$$

or

$$\langle\langle S_\rho u, U_j \rangle\rangle_{n-1} = \frac{1}{2}\pi \tilde{u}(j+1) + \sum_{k=2n-1-j}^{\infty} \tilde{u}(k) \langle\langle U_{k-1}, U_j \rangle\rangle_{n-1}, \quad 0 \leq j \leq n-2.$$

Similarly, there holds

$$\langle\langle S_\rho u_n, U_j \rangle\rangle_{n-1} = \frac{1}{2}\pi \tilde{u}_n(j+1), \quad 0 \leq j \leq n-2.$$

Using (15), we obtain

$$\tilde{u}_n(j+1) = \tilde{u}(j+1) + \frac{2}{\pi} \sum_{k=2n-j-1}^{\infty} \tilde{u}(k) \langle\langle U_{k-1}, U_j \rangle\rangle_{n-1}, \quad 0 \leq j \leq n-2. \quad (16)$$

To estimate the error  $u_n - u$  in the  $\tilde{H}_\rho^s$  norm we proceed as follows. Since  $\tilde{u}_n(k) = 0$  for  $k \geq n$ , we have

$$\|u_n - u\|_{\tilde{H}_\rho^s}^2 = \sum_{k=1}^{n-1} k^{2s} |\tilde{u}_n(k) - \tilde{u}(k)|^2 + \sum_{k=n}^{\infty} k^{2s} |\tilde{u}(k)|^2. \quad (17)$$

Using that Chebyshev polynomials satisfy [8] for  $j = 0, \dots, n-2$ ,

$$\langle\langle U_k, U_j \rangle\rangle_{n-1} = \begin{cases} -\frac{1}{2}\pi, & k = 2nm - 2 - j, \quad m = 1, 2, \dots, \\ \frac{1}{2}\pi, & k = 2nm + j, \quad m = 0, 1, \dots, \\ 0, & \text{otherwise,} \end{cases} \quad (18)$$

we estimate the second term in (17):

$$\sum_{k=n}^{\infty} k^{2s} |\tilde{u}(k)|^2 = \sum_{k=n}^{\infty} k^{2s-2t} k^{2t} |\tilde{u}(k)|^2 \leq n^{2s-2t} \|u\|_{\tilde{H}_\rho^t}^2,$$

provided  $s < t$ .

Finally, we consider the first term in (17). With (16) we have

$$\begin{aligned} & \sum_{j=0}^{n-2} (j+1)^{2s} |\tilde{u}_n(j+1) - \tilde{u}(j+1)|^2 \\ &= \sum_{j=0}^{n-2} (j+1)^{2s} \left( \frac{2}{\pi} \sum_{k=2n-j-1}^{\infty} \tilde{u}(k) \langle\langle U_{k-1}, U_j \rangle\rangle_{n-1} \right)^2 \\ &= \sum_{j=0}^{n-2} (j+1)^{2s} \left( \frac{2}{\pi} \sum_{k=2n-j-2}^{\infty} \tilde{u}(k+1) \langle\langle U_k, U_j \rangle\rangle_{n-1} \right)^2 \\ &= \sum_{j=1}^{n-2} (j+1)^{2s} \left( \sum_{m=1}^{\infty} \tilde{u}(2nm+j+1) - \tilde{u}(2nm-j-1) \right)^2, \quad \text{with (18).} \end{aligned}$$

If  $s \geq 0$ , we have  $(j+1)^{2s} \leq n^{2s}$ , whereas if  $s < 0$ , we have  $(j+1)^{2s} \leq 1$ . Thus for  $s \geq 0$  we obtain for  $t > \frac{1}{2}$ , using the Cauchy–Schwarz inequality,

$$\begin{aligned} & \sum_{j=0}^{n-2} (j+1)^{2s} |\tilde{u}_n(j+1) - \tilde{u}(j+1)|^2 \\ & \leq n^{2s} \sum_{j=0}^{n-2} \left| \sum_{m=1}^{\infty} \tilde{u}(2nm+j+1) - \sum_{m=1}^{\infty} \tilde{u}(2nm-j-1) \right|^2 \\ & \leq n^{2s} \sum_{j=0}^{n-2} \sum_{m=1}^{\infty} (2nm+j+1)^{-2t} \sum_{m=1}^{\infty} (2nm+j+1)^{2t} |\tilde{u}(2nm+j+1)|^2 \\ & \quad + n^{2s} \sum_{j=0}^{n-2} \sum_{m=1}^{\infty} (2nm-j-1)^{-2t} \sum_{m=1}^{\infty} (2nm-j-1)^{2t} |\tilde{u}(2nm-j-1)|^2 \\ & \leq Cn^{2s-2t} \sum_{j=0}^{n-2} \left[ \sum_{m=1}^{\infty} (2nm+j+1)^{2t} |\tilde{u}(2nm+j+1)|^2 \right. \\ & \quad \left. + \sum_{m=1}^{\infty} (2nm-j-1)^{2t} |\tilde{u}(2nm-j-1)|^2 \right] \\ & \leq Cn^{2s-2t} \sum_{k=n}^{\infty} k^{2t} |\tilde{u}(k)|^2 \leq Cn^{2s-2t} \|u\|_{\dot{H}_\rho^t}^2, \end{aligned}$$

using  $\sum_{m=1}^{\infty} (2nm-j-1)^{-2t} \leq \sum_{m=1}^{\infty} (nm)^{-2t} \leq Cn^{-2t}$ . For  $s < 0$  we estimate correspondingly, provided  $t > \frac{1}{2}$ .  $\square$

**Proof of Theorem 3.** Let  $f \in \dot{H}_{1/\rho}^s$  and  $S_\rho u = f$ . Then  $f(x) = \sum_{k=0}^{\infty} \hat{f}(k) U_k(x)$ , where

$$\|f\|_{\dot{H}_{1/\rho}^s}^2 = \sum_{k=0}^{\infty} |k+1|^{2s} |\hat{f}(k)|^2.$$

But due to (7) we have, with  $u = \sum_{k=0}^{\infty} \tilde{u}(k) T_k$  where  $\tilde{u}(0) = 0$  since  $u \in \tilde{H}_\rho^s \setminus \{T_0\}$ ,

$$S_\rho u = \sum_{k=0}^{\infty} \tilde{u}(k) S_\rho T_k = \sum_{k=1}^{\infty} \tilde{u}(k) U_{k-1} = \sum_{k=1}^{\infty} \hat{f}(k-1) U_{k-1},$$

yielding

$$\tilde{u}(k) = \hat{f}(k-1), \quad \text{for } k \geq 1.$$

Hence,

$$\|f\|_{\dot{H}_{1/\rho}^s}^2 = \sum_{k=0}^{\infty} |k+1|^{2s} |\tilde{u}(k+1)|^2 = \sum_{l=1}^{\infty} |l|^{2s} |\tilde{u}(l)|^2 = \|u\|_{\dot{H}_\rho^s}^2 - |\tilde{u}(0)|^2,$$

from which the assertion follows.  $\square$

**Remark 4.** For  $\tau = 0$  we have  $\hat{H}_{1/\rho}^\tau = L_{2,1/\rho}$ ,  $\tilde{H}_\rho^\tau = L_{2,\rho}$ , where

$$\|u\|_{L_{2,\rho}}^2 = \int_{-1}^1 \rho u^2 \, dx \quad \text{and} \quad \|f\|_{L_{2,1/\rho}}^2 = \int_{-1}^1 \frac{1}{\rho} f^2 \, dx.$$

The continuity of the mapping property  $S_\rho : L_{2,\rho} \rightarrow L_{2,1/\rho}$  is well known [9].

**Remark 5.** Due to Theorems 1 and 3 we have for  $f \in \hat{H}_{1/\rho}^t$  with  $t > s$  and  $t > \frac{1}{2}$  the estimate

$$\|u_n - u\|_{\tilde{H}_\rho^s} \leq \begin{cases} Cn^{-(t-s)} \|f\|_{\hat{H}_{1/\rho}^t}, & s \geq 0, \\ Cn^{-t} \|f\|_{\hat{H}_{1/\rho}^t}, & s < 0, \end{cases}$$

with the notation of Theorem 1.

## 2. Method B

For  $\alpha = \frac{1}{2}$ ,  $\beta = -\frac{1}{2}$  we look for solutions of (1) of the form

$$v(x) = \sqrt{\frac{1-x}{1+x}} u(x).$$

Due to property (6) it is natural to approximate the solution  $v$  of (1) by  $\rho(x) = (1-x)^{1/2}(1+x)^{-1/2}$  times a polynomial  $u_n$  of degree  $\leq n-1$ :

$$v_n(x) = \rho(x) \sum_{k=0}^{n-1} a_k p_k^{(1/2, -1/2)}(x), \quad a_k \in \mathbb{R}.$$

We choose the collocation points

$$x_k = \cos \frac{2k-1}{2n+1} \pi, \quad k = 1, 2, \dots, n, \quad (19)$$

which are the distinct zeros of  $p_n^{(-1/2, 1/2)}(x)$ . Thus our collocation scheme is: find  $u_n \in \mathbb{P}_{n-1}$  such that

$$S_\rho u_n(x_k) = f(x_k), \quad 1 \leq k \leq n. \quad (20)$$

Here,

$$\begin{aligned} p_n^{(-1/2, 1/2)}(x) &= \frac{1}{\sqrt{\pi}} \frac{\cos((n + \frac{1}{2})\cos^{-1}x)}{\cos(\frac{1}{2}\cos^{-1}x)}, \quad n = 0, 1, \dots, \\ p_n^{(1/2, -1/2)}(x) &= \frac{1}{\sqrt{\pi}} \frac{\sin((n + \frac{1}{2})\cos^{-1}x)}{\sin(\frac{1}{2}\cos^{-1}x)}, \quad n = 0, 1, \dots. \end{aligned} \quad (21)$$

Now, with the aid of (6),

$$S_\rho u_n(x) = - \sum_{k=0}^{n-1} a_k p_k^{(-1/2, 1/2)}(x). \quad (22)$$



Note that (20) represents a set of linear equations for  $a_0, \dots, a_{n-1}$  with easily computed matrix elements. Again, one may even obtain explicit expressions for  $a_0, \dots, a_{n-1}$  by exploiting the discrete orthogonality of the Jacobi polynomials  $\{p_k^{(-1/2, 1/2)}(x)\}_{k=0}^\infty$ . There holds the following discrete inner product (obtained by using the Gauss–Jacobi quadrature rule based on the weight  $(1/\rho)(x) = (1-x)^{-1/2}(1+x)^{1/2}$  which uses the zeros (19) of  $p_n^{(-1/2, 1/2)}(x)$  as nodes and the weights  $\omega_k$  (see [12])):

$$\langle u, w \rangle_n = \sum_{k=1}^n \omega_k u(x_k) w(x_k). \quad (23)$$

Then there holds

$$\begin{aligned} \langle p_j^{(-1/2, 1/2)}, p_k^{(-1/2, 1/2)} \rangle_n &= \int_{-1}^1 p_j^{(-1/2, 1/2)}(x) p_k^{(-1/2, 1/2)}(x) \frac{1}{\rho}(x) dx \\ &= \delta_{jk}, \quad j, k \leq n-1. \end{aligned}$$

Hence, from (22) follows

$$a_k = -\langle f, p_k^{(-1/2, 1/2)} \rangle_n, \quad k = 0, \dots, n-1.$$

With

$$\begin{aligned} \|u\|_{\bar{H}_\rho^s}^2 &= \sum_{k=1}^\infty |k|^{2s} |\hat{u}(k)|^2 + |\hat{u}(0)|^2, \\ u(x) &= \sum_{k=0}^\infty \hat{u}(k) p_k^{(1/2, -1/2)}(x), \\ \hat{u}(k) &= \int_{-1}^1 u(x) p_k^{(1/2, -1/2)}(x) \rho(x) dx, \end{aligned} \quad (24)$$

we have the following theorem.

**Theorem 6.** Let  $u \in \bar{H}_\rho^t$  solve (1) with  $u = v/\rho$  and  $\rho(x) = (1-x)^{1/2}(1+x)^{-1/2}$ . Then for any  $n \in \mathbb{Z}_+$  there exists a solution  $u_n \in \mathbb{P}_{n-1}$  of (20). Furthermore, if  $t > s$  and  $t > \frac{1}{2}$ , then  $u_n$  satisfies

$$\|u_n - u\|_{\bar{H}_\rho^s} \leq \begin{cases} Cn^{-(t-s)} \|u\|_{\bar{H}_\rho^t}, & s \geq 0, \\ Cn^{-t} \|u\|_{\bar{H}_\rho^t}, & s < 0, \end{cases} \quad (25)$$

with  $C$  a constant independent of  $n$ .

**Proof.** The approximation scheme (20) is equivalent to

$$\langle S_\rho u_n, p_j^{(-1/2, 1/2)} \rangle_n = \langle S_\rho u, p_j^{(-1/2, 1/2)} \rangle_n, \quad j = 0, \dots, n-1. \quad (26)$$

Again one needs a discrete orthogonality property like (18) but now for the orthogonal system  $\{p_k^{(-1/2, 1/2)}(x)\}_{k=0}^\infty$  with respect to the discrete inner product in (23). This discrete orthogonality

property can be obtained by using (21). One obtains for  $j = 0, \dots, n-1$ ,

$$\langle p_k^{(-1/2, 1/2)}, p_j^{(-1/2, 1/2)} \rangle_n = \begin{cases} -(-1)^m, & k = (2n+1)m - 1 - j, \quad m = 1, 2, \dots, \\ (-1)^m, & k = (2n+1)m + j, \quad m = 0, 1, \dots, \\ 0, & \text{otherwise.} \end{cases} \quad (27)$$

Now from (6) and (24) with  $v = \rho u$ ,  $\rho = \sqrt{(1-x)/(1+x)}$  we have

$$S_\rho u(x) = - \sum_{k=0}^{\infty} \hat{u}(k) p_k^{(-1/2, 1/2)}(x).$$

Thus we can compute with (27) for  $j = 0, \dots, n-1$ ,

$$\begin{aligned} \langle S_\rho u, p_j^{(-1/2, 1/2)} \rangle_n &= - \sum_{k=0}^{\infty} \hat{u}(k) \langle p_k^{(-1/2, 1/2)}, p_j^{(-1/2, 1/2)} \rangle_n \\ &= - \sum_{m=0}^{\infty} \sum_{r=0}^{2n} \hat{u}(r + (2n+1)m) \langle p_{(2n+1)m+r}^{(-1/2, 1/2)}, p_j^{(-1/2, 1/2)} \rangle_n \\ &= - \sum_{m=0}^{\infty} (-1)^m \hat{u}(j + (2n+1)m) + \sum_{m=1}^{\infty} (-1)^m \hat{u}(-j-1 + (2n+1)m) \\ &= -\hat{u}(j) - \sum_{m=1}^{\infty} (-1)^m \hat{u}(j + (2n+1)m) + \sum_{m=1}^{\infty} (-1)^m \hat{u}(-j-1 + (2n+1)m). \end{aligned} \quad (28)$$

On the other hand, there holds for  $j = 0, \dots, n-1$ ,

$$\langle S_\rho u_n, p_j^{(-1/2, 1/2)} \rangle_n = -\hat{u}_n(j). \quad (29)$$

Thus one obtains from (26) with (28) and (29) for  $j = 0, \dots, n-1$ ,

$$\hat{u}(j) - \hat{u}_n(j) = - \sum_{m=1}^{\infty} (-1)^m \hat{u}(j + (2n+1)m) + \sum_{m=1}^{\infty} (-1)^m \hat{u}(-j-1 + (2n+1)m).$$

Next we estimate the norm in (24):

$$\|u_n - u\|_{\tilde{H}_\rho^s}^2 = \sum_{j=0}^{n-1} j^{2s} |\hat{u}_n(j) - \hat{u}(j)|^2 + \sum_{j=n}^{\infty} j^{2s} |\hat{u}(j)|^2. \quad (30)$$

First we estimate with the second term in (30) for  $s < t$ ,

$$\sum_{j=n}^{\infty} j^{2s} |\hat{u}(j)|^2 = \sum_{j=n}^{\infty} j^{2s-2t} j^{2t} |\hat{u}(j)|^2 \leq n^{2s-2t} \|u\|_{\tilde{H}_\rho^t}^2.$$

Next we consider with the first term in (30) for  $s \geq 0$  and  $t > s$ ,

$$\begin{aligned} & \left| \sum_{j=0}^{n-1} j^{2s} \left[ - \sum_{m=1}^{\infty} (-1)^m \hat{u}(j + (2n+1)m) + \sum_{m=1}^{\infty} (-1)^m \hat{u}(-j-1 + (2n+1)m) \right] \right|^2 \\ & \leq n^{2s} \sum_{j=0}^{n-1} \left( \sum_{m=1}^{\infty} (j + (2n+1)m)^{-2t} \sum_{m=1}^{\infty} (j + (2n+1)m)^{2t} |\hat{u}(j + (2n+1)m)|^2 \right. \\ & \quad \left. + \sum_{m=1}^{\infty} (-j-1 + (2n+1)m)^{-2t} \sum_{m=1}^{\infty} (-j-1 + (2n+1)m)^{2t} |\hat{u}(-j-1 + (2n+1)m)|^2 \right) \\ & \leq C n^{2s-2t} \sum_{k=n+1}^{\infty} k^{2t} |\hat{u}(k)|^2 \leq C n^{2s-2t} \|u\|_{\dot{H}_p^t}^2. \end{aligned}$$

For  $s < 0$  we use  $j^{2s} \leq 1$  and obtain analogously the desired estimate (25), provided  $t > \frac{1}{2}$ .  $\square$

### 3. Numerical results

The following numerical results were performed on the CDC 855 at the University of Hannover.

Table 1  
Weighted  $L^2$ -error for collocation scheme (9)

$(1-x^2)^{3/2}$			$e^{-x}$		
$n$	$\ e\ _{\dot{H}_p^0}$	$\alpha_n$	$n$	$\ e\ _{\dot{H}_p^0}$	$\alpha_n$
6	$0.525 \cdot 10^{-2}$	4.56	3	0.170	6.31
11	$0.332 \cdot 10^{-3}$	5.15	4	$0.277 \cdot 10^{-1}$	9.38
16	$0.481 \cdot 10^{-4}$	4.41	5	$0.342 \cdot 10^{-2}$	12.68
21	$0.145 \cdot 10^{-4}$	4.85	6	$0.339 \cdot 10^{-3}$	16.16
26	$0.515 \cdot 10^{-5}$	4.46	7	$0.281 \cdot 10^{-4}$	19.80
31	$0.235 \cdot 10^{-5}$	4.79	8	$0.199 \cdot 10^{-5}$	23.57
36	$0.115 \cdot 10^{-5}$	4.51	9	$0.124 \cdot 10^{-6}$	27.46
41	$0.638 \cdot 10^{-6}$	4.78	10	$0.688 \cdot 10^{-8}$	31.46
46	$0.368 \cdot 10^{-6}$	4.57	11	$0.343 \cdot 10^{-9}$	35.55
51	$0.230 \cdot 10^{-6}$		12	$0.156 \cdot 10^{-10}$	
$\alpha^*$		4.5			

In Table 1 we give the weighted  $L^2$ -error for the collocation approximation of (1) and present the associated experimental convergence rates  $\alpha_n$ .

For  $f$  listed in Table 1 we approximate the unknown solution  $v$  of (1) by taking  $\tilde{v} = \rho u_{200}^*$  with  $\rho(x) = 1/\sqrt{1-x^2}$  as proposed in Method A. Here  $u_{200}^*$  solves the collocation scheme (9) for  $n = 200$ . Thus instead of the actual error  $u - u_n$ , where  $u = v/\rho$  and  $u_n$  solves (9) and satisfies the additional constraint (11), we list in Table 1 the weighted  $L^2$ -error for  $e = u_{200}^* - u_n$ , i.e.,

$$\|e\|_{\tilde{H}_\rho^0} = \left( \int_{-1}^1 (1-x^2)^{-1/2} |u_{200}^* - u_n|^2 dx \right)^{1/2}. \quad (31)$$

In evaluating (31) we use the respective Chebyshev polynomial expansions (8) for  $u_{200}^*$  and  $u_n$  and perform the integration by the 200-point Gauss–Chebyshev quadrature rule. Note that  $(1-x^2)^{3/2} \in \hat{H}_{1/\rho}^{9/2-\epsilon}$  for any  $\epsilon > 0$ , and  $e^{-x} \in \hat{H}_{1/\rho}^s$  for any  $s$ , which can be seen by computing the Chebyshev coefficients [7]. Therefore Theorem 1 together with Theorem 3 gives the theoretical convergence rate  $\alpha^*$  listed in Table 1 (cf. Remark 5). For  $f = e^{-x}$  Table 1 shows faster than polynomial convergence.

## Acknowledgement

The authors would like to thank Prof. Dr. D. Elliott for valuable remarks.

## References

- [1] M.J. Dow and D. Elliott, The numerical solution of singular integral equations over  $(-1, 1)$ , *SIAM J. Numer. Anal.* **16** (1979) 115–134.
- [2] A.V. Dzishkariani, The solution of singular integral equations by approximate projection methods, *U.S.S.R. Comput. Math. and Math. Phys.* **19** (1980) 61–74.
- [3] D. Elliott, Orthogonal polynomials associated with singular integral equations having a Cauchy kernel, *SIAM J. Math. Anal.* **13** (1982) 1041–1052.
- [4] D. Elliott, The classical collocation method for singular integral equations, *SIAM J. Numer. Anal.* **19** (1982) 816–832.
- [5] D. Elliott, Rates of convergence for the method of classical collocation for solving singular integral equations, *SIAM J. Numer. Anal.* **21** (1984) 136–148.
- [6] D. Elliott, A comprehensive approach to the approximate solution of singular integral equations over the area  $(-1, 1)$ , *J. Integral Equations Appl.* **2** (1989) 59–91.
- [7] V.J. Ervin and E.P. Stephan, Collocation with Chebyshev polynomials for first kind integral equations on intervals, in: J.R. Whiteman, Ed., *The Mathematics of Finite Elements and Applications VII* (Academic Press, New York, 1991) 321–329.
- [8] V.J. Ervin and E.P. Stephan, Collocation with Chebyshev polynomials for a hypersingular integral equation on an interval, *J. Comput. Appl. Math.* **43** (1–2) (1992) 221–229.
- [9] M.A. Golberg, The convergence of a collocation method for a class of Cauchy singular integral equations, *J. Math. Anal. Appl.* **100** (1984) 500–512.
- [10] M.A. Golberg and J.A. Fromme, On the  $L^2$  convergence of collocation for the generalized airfoil equation, *J. Math. Anal. Appl.* **71** (1979) 271–286.

- [11] P. Junghanns and B. Silbermann, Theorie der Näherungsverfahren für singuläre Integralgleichungen auf Intervallen, *Math. Nachr.* **103** (1981) 199–244.
- [12] S. Krenk, On quadrature formulas for singular integral equations of the first and the second kind, *Quart. Appl. Math.* **33** (1975) 225–232.
- [13] I.H. Sloan and E.P. Stephan, Collocation with Chebyshev polynomials for Symm's integral equation on an interval, *J. Austral. Math. Soc. Ser. B* **34** (1992) 199–211.